

vector passing through two points (x_1, y_1, z_1) and (x_2, y_2, z_2) :

Position vector of P is

$$\vec{OP} = \vec{r}_1 = x_1 \hat{i} + y_1 \hat{j} + z_1 \hat{k}$$

Position vector of A is

$$\vec{OA} = \vec{r}_2 = x_2 \hat{i} + y_2 \hat{j} + z_2 \hat{k}$$


When in $\triangle OPA$,

$$\vec{OP} + \vec{PA} = \vec{OA}$$


$$\therefore \vec{PA} = \vec{OA} - \vec{OP}$$

$$= (x_2 \hat{i} + y_2 \hat{j} + z_2 \hat{k}) - (x_1 \hat{i} + y_1 \hat{j} + z_1 \hat{k})$$

$$= (x_2 - x_1) \hat{i} + (y_2 - y_1) \hat{j} + (z_2 - z_1) \hat{k}$$



Exp:



$$\text{When } \vec{PQ} = (5-2) \hat{i} + (3-4) \hat{j} + (4-3) \hat{k}$$

$$= 3 \hat{i} - \hat{j} + \hat{k}$$


Projection of a vector on other vector:

Let $\vec{A} = a_1 \hat{i} + a_2 \hat{j} + a_3 \hat{k}$ is a given vector.

$\vec{B} = b_1 \hat{i} + b_2 \hat{j} + b_3 \hat{k}$ is another vector.

When unit vector in the direction of vector \vec{B} is \hat{b} (suppose).

Now projection of the vector \vec{A} on the other vector \vec{B} is $\vec{A} \cdot \hat{b}$.



Problems:-

Q1: Find the values of a , when the vectors $\vec{A} = a\hat{i} - \hat{j} + \hat{k}$ and $\vec{B} = 2a\hat{i} + a\hat{j} - 4\hat{k}$ are perpendicular.

Ans:- $\vec{A} = a\hat{i} - \hat{j} + \hat{k}$
 $\vec{B} = 2a\hat{i} + a\hat{j} - 4\hat{k}$.

$$\vec{A} \cdot \vec{B} = |\vec{A}| |\vec{B}| \cos 90^\circ$$

Since the two vectors are perpendicular to each other,

$$\therefore \vec{A} \cdot \vec{B} = 0$$

$$(a\hat{i} - \hat{j} + \hat{k}) \cdot (2a\hat{i} + a\hat{j} - 4\hat{k}) = 0$$

$$2a^2 - 2a - 4 = 0$$

$$a^2 - a - 2 = 0$$

$$(a-2)(a+1) = 0$$


$$\therefore a = \underline{\underline{2, -1}}$$

Q2:- Find the projection of the vector $2\hat{i} - 3\hat{j} + 6\hat{k}$ on the vector $\hat{i} + 2\hat{j} + 2\hat{k}$.

Ans:- let $\vec{A} = 2\hat{i} - 3\hat{j} + 6\hat{k}$; and
 $\vec{B} = \hat{i} + 2\hat{j} + 2\hat{k}$.

Unit vector in the direction of \vec{B} is $\hat{b} = \frac{\hat{i} + 2\hat{j} + 2\hat{k}}{\sqrt{1^2 + 2^2 + 2^2}}$

$$= \frac{1}{3} (\hat{i} + 2\hat{j} + 2\hat{k})$$



Now projection of \vec{A} on \vec{B} is $= \vec{A} \cdot \hat{b}$

$$= (2\hat{i} - 3\hat{j} + 6\hat{k}) \cdot \frac{1}{3} (\hat{i} + 2\hat{j} + 2\hat{k})$$


$$= \frac{1}{3} (2 - 6 + 12)$$

$$= \frac{8}{3} \text{ Ans:}$$

Q3: Find the work done in moving an object along the straight line from $(3, 2, -1)$ to $(2, -1, 4)$ in a force field given by $\vec{F} = 4\hat{i} - 3\hat{j} + 2\hat{k}$.

Ans:- Let \vec{r} be the vector passes through the points from $(3, 2, -1)$ to $(2, -1, 4)$. Then

$$\begin{aligned}\vec{r} &= (2-3)\hat{i} + (-1-2)\hat{j} + (4+1)\hat{k} \\ &= -\hat{i} - 3\hat{j} + 5\hat{k}\end{aligned}$$




Now work done = (magnitude of force in the direction of motion) (distance moved)

$$\begin{aligned}&= (F \cos \theta) (r) \\ &= F r \cos \theta = \vec{F} \cdot \vec{r} \\ &= (4\hat{i} - 3\hat{j} + 2\hat{k}) \cdot (-\hat{i} - 3\hat{j} + 5\hat{k}) \\ &= -4 + 9 + 10 = 15 \text{ Ans.}\end{aligned}$$

Q4: Find the volume of the parallelepiped

Note: Cross product (vector product):



$$\vec{A} \times \vec{B} = |\vec{A}| |\vec{B}| \sin \theta \hat{n}$$

Note: Let $\vec{A} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$
 $\vec{B} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$

then

$$\vec{A} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$